

This article was downloaded by: [Barnes, Maria E.]

On: 17 December 2009

Access details: Access Details: [subscription number 914693504]

Publisher Psychology Press

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Developmental Neuropsychology

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t775653638>

### Impairments in Attention in Occasionally Snoring Children: An Event-Related Potential Study

Maria E. Barnes <sup>a</sup>; Elizabeth A. Huss <sup>a</sup>; Krista N. Garrod <sup>a</sup>; Eric Van Raay <sup>a</sup>; Ehab Dayyat <sup>b</sup>; David Gozal <sup>c</sup>; Dennis L. Molfese <sup>d</sup>

<sup>a</sup> Department of Psychological and Brain Science, University of Louisville, Louisville, Kentucky <sup>b</sup>

Department of Pediatrics, University of Louisville, Louisville, Kentucky <sup>c</sup> Department of Pediatrics,

Pritzker School of Medicine, Comer Children's Hospital, University of Chicago, Chicago, Illinois <sup>d</sup> Birth

Defects Center, University of Louisville, Louisville, Kentucky

**To cite this Article** Barnes, Maria E., Huss, Elizabeth A., Garrod, Krista N., Van Raay, Eric, Dayyat, Ehab, Gozal, David and Molfese, Dennis L.(2009) 'Impairments in Attention in Occasionally Snoring Children: An Event-Related Potential Study', *Developmental Neuropsychology*, 34: 5, 629 — 649

**To link to this Article:** DOI: 10.1080/87565640903133632

URL: <http://dx.doi.org/10.1080/87565640903133632>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## Impairments in Attention in Occasionally Snoring Children: An Event-Related Potential Study

Maria E. Barnes, Elizabeth A. Huss, Krista N. Garrod, and Eric Van Raay  
*Department of Psychological and Brain Science*  
*University of Louisville, Louisville, Kentucky*

Ehab Dayyat  
*Department of Pediatrics*  
*University of Louisville, Louisville, Kentucky*

David Gozal  
*Department of Pediatrics, Pritzker School of Medicine, Comer Children's Hospital,*  
*University of Chicago, Chicago, Illinois*

Dennis L. Molfese  
*Birth Defects Center*  
*University of Louisville, Louisville, Kentucky*

**Objective:** To determine whether minimal snoring is benign in children. **Procedure:** 22 rarely snoring children (mean age = 6.9 years, 11 females) and age- and sex-matched controls participated in an auditory oddball task wearing 128-electrode nets. Parents completed the Conners Parent Rating Scales–Revised Long (CPRS–R:L). **Results:** Snorers scored significantly higher on four CPRS–R:L subscales. Stepwise regression indicated that two ERP variables from a region of the ERP that peaked at 844 msec post-stimulus onset predicted CPRS–R:L Attention Deficit Hyperactivity Disorder (ADHD) Index scores. **Conclusions:** Occasional snorers, according to parental report, do exhibit ADHD-like behaviors. Basic sensory processing is longer than in controls, suggesting that delayed frontal activation requires more effort in snorers.

Over the past decade, the impact of sleep-disordered breathing (SDB) in children on cognitive functions has received substantial attention (Arman et al., 2005; O'Brien, Tauman, & D. Gozal, 2004; Rosen et al., 2004; Urschitz et al., 2003). SDB has been repeatedly associated with multiple behavioral and neurocognitive dysfunctions, including restlessness, aggression, excessive daytime sleepiness, and poor school performance (Adams, Schluchter, & Redline, 2001; Ali, Pitson, & Stradling,

---

Correspondence should be addressed to Maria E. Barnes, Department of Psychological and Brain Sciences, University of Louisville, Louisville, KY 40292. E-mail: mebarn02@louisville.edu

Research was supported by a dissertation research award from the American Psychological Association; a fellowship from the National Institute of Mental Health (F30MH79531), and the following grants from the National Institutes of Health: R01HL070911, R01HL65270, R01DC005994, and R41HD47083. The experiments described herein were conducted as part of a larger, federally funded research study. Portions of this article have been presented in poster format at a national scientific conference.

1996; Beebe & Gozal, 2002; Gottlieb et al., 2003; Gozal, 1998; D. Gozal, Holbrook, Mehl, et al., 2001; D. Gozal & Pope, 2001; Hill, 1889; Leach, Olson, Hermann, & Manning, 1992; Rhodes et al., 1995; Singer, 1990; Weissbluth, Poncher, & Reiff, 1983). In addition, an association between SDB and attention deficit hyperactivity disorder (ADHD) in children has been documented by multiple investigators (Ball, Tiernan, Janusz, & Furr, 1997; Corkum, Tannock, Moldofsky, Hogg-Johnson, & Humphries, 2001; Kaplan, McNichol, Conte, & Moghadam, 1987; Owens, Maxim, Nobile, McGuinn, & Msall, 2000; Ring et al., 1998; Stein, 1999). The need for early diagnosis and treatment of pediatric SDB is further highlighted by the recognition that the age at which SDB is most prevalent (1–8 years) coincides with an important period for neuropsychological development in both humans and animals (Beebe & Gozal, 2002; D. Gozal, Daniel, & Dohanich, 2001; E. Gozal, Row, Schurr, & Gozal, D., 2001). As further testament of the adverse long-term effects of SDB, Gozal and Pope reported that children who snored during early childhood were at greater risk for poor academic performance in later years, long after snoring had resolved (D. Gozal & Pope, 2001). This unique vulnerability to insult by SDB in the rapidly developing brain could be a key difference in adult and pediatric forms of the disease (Halbower & Mahone, 2006; E. Gozal et al., 2001).

In 2002, the American Academy of Pediatrics (AAP) recommended that all children who habitually snore should be evaluated (American Academy of Pediatrics, 2002), while earlier literature suggested that the presence of snoring was important only insofar as it served as a potential, albeit unreliable, marker for more severe forms of SDB such as obstructive sleep apnea (OSA) (Ali, Pitson, & Stradling, 1993; Gislason & Benediktsdottir, 1995; D. Gozal & Pope, 2001; Guilleminault, Eldridge, Simmons, & Dement, 1976; Guilleminault, Winkle, Korobkin, & Simmons, 1982). Based on recent studies reporting neurocognitive dysfunction and impaired daytime performance in children who snore in the absence of overt or clinically diagnosed OSA (Alexopoulos et al., 2006; Guilleminault & Lee, 2004; Lopes & Guilleminault, 2006; O'Brien, Mervis, Holbrook, Bruner, Klaus, Rutherford, et al., 2004a; O'Brien, Mervis, Holbrook, Bruner, Smith, McNally, et al., 2004), some authors proposed that parental report of snoring may operate as one of the most robust predictors of cognition and school performance (Beebe, 2006). Habitually snoring children are at higher risk for impaired academic performance, increased social problems, decreased attention, impaired visuospatial ability, and poorer scores on measures of anxiety and depression (Blunden, Lushington, Lorenzen, Martin, & Kennedy, 2005; Chervin, Ruzicka, Archbold, & Dillon, 2005; O'Brien, Mervis, Holbrook, Bruner, Klaus, Rutherford, et al., 2004a; Urschitz et al., 2004; Urschitz et al., 2003). Consequently, habitual snoring may soon become considered as the mildest form of SDB, rather than be dismissed as clinically irrelevant (Guilleminault & Lee, 2004). The case for occasional snoring in non-habitual snorers having no adverse consequence has been largely ignored in the literature under the assumption that occasional snoring should be viewed as a truly benign condition (Kuehni, Strippoli, Chauliac, & Silverman, 2008; Urschitz et al., 2004). Unfortunately, occasional snorers have not been considered a distinct group, and their inclusion in studies investigating SDB in children has been rather inconsistent. Most often, these non-habitual snorers are combined with non-snorers in the control group of children who do not have SDB (Karpinski, Scullin, & Montgomery-Downs, 2008; Liukkonen, Virkkula, Aronen, Kirjavainen, & Pitkaranta, 2008; O'Brien, Mervis, Holbrook, Bruner, Klaus, Rutherford 2004a; Urschitz et al., 2003). It is possible that these methodological inconsistencies and the underlying assumption that occasional snoring imposes no adverse consequences may account for the high degree of variability across reports concerning the neurobehavioral sequelae associated with SDB in children.

To examine whether a putatively benign condition such as occasional snoring is associated with altered brain function, we applied a highly sensitive approach, namely event related potentials (ERPs). This method—and the attention paradigm described later in this article—has been used in adults with SDB to investigate patterns of brain activation underlying behaviorally observed deficits in domains such as attention (Gosselin et al., 2006a; Gosselin et al., 2006b). The recording of ERPs is a highly sensitive technique that is both safe and well tolerated in children and infants. ERPs can detect (and in some cases predict) significant neuropsychological differences even in the absence of gross findings (Lyytinen et al., 2001; Lyytinen, Blomberg, & Naatanen, 1992; Lyytinen & Naatanen, 1987; Molfese & Molfese, 1985; Molfese, 1989, 1990, 2000; Molfese et al., 2006).

We hypothesized that occasional snoring would disrupt neuropsychological functioning with snoring children exhibiting impaired activation over frontal and prefrontal areas, areas normally associated with attention. The experiments described herein examined differences in ERPs recorded during a classic attention task between a group of pre-pubertal community children who snored “rarely” or “occasionally” according to parental report and a matched control group who had no history of snoring. Dependent ERP variables were then analyzed in relation to sleep measures and scores on a standardized and sensitive measure of neurobehavioral functioning in children with SDB, the Conners Parent Rating Scales–Revised: Long (CPRS–R:L) (Wei, Mayo, Smith, Reese, & Weatherly, 2007). Furthermore, this decrease in cognitive reserve was hypothesized to predict observable neurobehavioral problems as measured by the CPRS–R:L.

## METHODS

### Participants

Participants included 44 children between 4 and 8 years of age (mean age = 6.92 years,  $SD = 1.24$ , 22 females). Subjects were selected on the basis of parental responses to items on the Sleep Behavior Questionnaire (SBQ, described later) concerning snoring frequency and severity. Children who were reported to snore either “Rarely” or “Occasionally,” were assigned to the non-habitual snoring group. Children for whom parents selected “Never” were assigned to the control group. For both groups, the presence or absence of snoring was confirmed during an overnight sleep study (PSG). Children were tested as part a larger, ongoing study of sleep characteristics in healthy children. They were financially compensated along with a bag of books, puzzles, and toys for their participation in the larger study. All subjects were recruited and consented according to the guidelines of the American Psychological Association. Approval of the Institutional Review Board of the University of Louisville was obtained prior to the start of the study. A photograph of a participant wearing the 128-electrode net used to record the ERPs is shown in Figure 1.

Exclusion criteria for both groups included: pre-existing medical, neurological, psychiatric, behavioral, or learning disorders indicated by screening questionnaires on history; medication usage; obesity (BMI greater than 95th percentile); significant growth failure; developmental delay; failure to pass the hearing and vision screening (using air conduction threshold tests in the same normal range for both ears as a hearing test and tumbling-E Snellen charts for a measure of approximate visual acuity); failure to pass IQ screening (using the Peabody Picture Vocabulary Test III as a proxy for IQ described below); first-degree relative with a major affective disorder or mental illness assessed by family history diagnostic criteria; and significant “at-risk” scores on the



FIGURE 1 Photograph of participant in 128-electrode net.

Childhood Symptom Inventory-4 (described later). A licensed psychologist clinically interviewed children (and their parents) whom the CSI-4 determined to be "at risk." Children with significant behavioral disorders were excluded. The final sample included 22 children in the non-habitual snoring group and 22 age- and sex-matched controls. Both groups comprised 11 males and 11 females each. Table 1 outlines pertinent demographic information for this sample. There were no significant group differences for any demographic variable.

## Apparatus and Procedure

### *ERP Apparatus and Procedures*

**Equipment.** A high-density 128 Ag/AgCl electrode system utilizing high-impedance amplifiers (EGI, Eugene, OR) was used to record ERP activity. NetStation © software (EGI, Eugene, OR) controlled impedance measures, baseline correction, analog/digital sampling rate, artifact rejection, and averaging. A second computer used e-prime © software to present stimuli, integrating this information with the ERP recording software. Interstimulus intervals (ISI) varied randomly between all stimuli.

**Testing methods.** Children were tested individually in sound-dampened rooms at the Sleep Center the morning following their overnight PSG sleep study. Screening tests were administered, including hearing (air conduction threshold) and vision screening (Snellen charts) and behavioral testing. The child's head was measured, and the vertex was marked for proper alignment of electrode net placement. Following standard procedures, the electrode net was soaked for 5 minutes in warm KCl solution. Net application and impedance adjustments required less than 5 minutes. Electrode impedances, recorded prior to and at the end of each task, were below 40 kOhms to maximize signal-to-noise ratio, thereby produce a high-quality signal for subsequent analyses. ERP signals were digitized online at 4 msec intervals for a 1.5 sec period for each of the 128 sites. Filters were set at 0.1 Hz for high pass and 30 Hz for low pass, with a gain of 10 K. The system col-

TABLE 1  
Demographic and Behavioral Variables for Rarely Snoring  
and Non-Snoring Children

Variable	Snoring Group	Control Group
Age	6.9195 ± 1.2443	6.9186 ± 1.2069
BMI	17.5628 ± 4.1205	15.9886 ± 1.3172
PPVT-III	108.5882 ± 12.3999	106.4667 ± 13.0541
NEPSY Visual Attention	10.9545 ± 3.1088	10.5714 ± 3.2950

Both groups have  $N = 22$ . Values given in means ± SD. PPVT-III = Standardized score on Peabody Picture Vocabulary Test-III. NEPSY Visual Attention = Scaled score on NEPSY Visual Attention subtest.

lected auditory and visual ERPs as the child participated in 3 tasks. If the child moved, recording was paused until the child was sitting quietly. Children took 3-min breaks at the end of each task. ERP testing lasted about 50 min.

**Electrophysiological task.** The “oddball” or “P300” task is a classic attention task in the ERP field (Sutton, Tueting, Zubin, & John, 1965). The findings have been consistent across numerous studies over a variety of populations, including studies of sleep apnea. In fact, the P300 peak (also referred to as subtype P3b) is the most widely studied ERP component in adults and children. The results of this task are related to a wide range of behavioral assessments of attention and provide a direct link to the cognitions of attention and vigilance. The amplitude of a positive peak occurring about 300 msec after stimulus onset is larger in response to infrequent events in adults. In healthy children, this peak typically occurs approximately 400 msec (Emmerson, Dustman, Shearer, & Turner, 1990; Key, Dove, & Maguire, 2005; Polich & Martin, 1992).

**Stimuli.** Two single frequency tones, 300 msec duration, selected from a range between 1000 and 4000 Hz served as stimuli. Tones presented during any test differed by 500 Hz. Rise and decay times were matched between stimulus pairs. Auditory stimuli had matched loudness levels, 75 dB SPL(A), and were presented through a speaker one meter over the midline of the child’s head.

**Pre-test training.** Before testing, the child listened to a series of two “beeps” that occurred at 1-sec intervals. One tone occurred with high frequency (four times in a row). The second (target) always occurred on the 5th, 10th, 15th, and 20th trials. Children quickly learned to press the correct button to the tone presented in less than 20 trials.

**Task.** The experimental task presented two tones at 2-sec intervals. Tones occurred in random order, so that one occurred on 70% of the trials while the other occurred on 30%. To decrease habituation and learning, tone frequency changed across test sessions for each child, and frequency of occurrence for the higher tone versus the lower was counterbalanced across sessions, children, and sex. The test presented 120 tones. The tester asked the child to attend to the target and press a button on hearing it. Tone presentation lasted 5 min.



### *PSG Apparatus and Procedures*

All children and one of their parents reported to the Sleep Laboratory around 7:30 pm. Following adaptation to the testing environment and personnel, the sleep studies began between 9:00 and 9:30 pm. The following parameters were measured: chest and abdominal wall movement by respiratory impedance or inductance plethysmography, heart rate by electrocardiogram (ECG); air flow with a sidestream end-tidal capnograph (PETCO<sub>2</sub>; BCI SC-300, Menomonee Falls, WI) which also provided breath-by-breath assessment of end-tidal carbon dioxide levels, nasal pressure and an oral-nasal airflow thermistor. Arterial oxygen saturation (SpO<sub>2</sub>) was assessed by pulse oximetry (Nellcor N 100; Nellcor Inc., Hayward, CA), with simultaneously recorded pulse waveform. Bilateral electro-oculogram (EOG), 8 channels of electroencephalogram (EEG), chin and anterior tibial electromyograms (EMG), and analog output from a body position sensor (Braebon Medical Corporation, NY) were also monitored. All measures were digitized using a commercially available computerized PSG system (MedCare Diagnostics, Buffalo, NY or Stellate Systems, Montreal, Quebec, Canada). Tracheal sound was monitored with a microphone sensor (Sleepmate, VA) for detection of snoring and digital time-synchronized video images were collected. The sleep technician followed patient behavior and confirmed sleep position by the infrared camera inside the room. The presence of snoring was verified in occasionally snoring children provided that at least 10 min of observed snoring would be present during sleep. Conversely, the absence of snoring was confirmed as well for control children. All the studies were initially scored for both sleep stages and respiratory events by a certified technician and were then reviewed by physicians experienced in pediatric polysomnography (Rechtschaffen & Kales, 1968). The proportion of time spent in each sleep stage was expressed as percentage of total sleep time (%TST). Central, obstructive, and mixed apneic events were counted. Obstructive apnea was defined as the absence of airflow with continued chest wall and abdominal movement for duration of at least two breaths (Montgomery-Downs, O'Brien, Gulliver, & Gozal, 2006). Hypopneas were defined as a decrease in oronasal flow of  $\geq 50\%$  with a corresponding decrease in SpO<sub>2</sub> of  $\geq 4\%$  and/or arousal (Montgomery-Downs et al., 2006). The obstructive apnea/hypopnea index (AHI) was defined as the number of apneas and hypopneas per hour of TST. The mean oxygen saturation, as measured by pulse oximetry (SpO<sub>2</sub>) in the presence of a pulse waveform signal void of motion artifact, and the nadir SpO<sub>2</sub> were recorded. Arousals were expressed as the total number of arousals per hour of sleep time (arousal index), and were defined as recommended by the (American Sleep Disorders Association Task Force, 1992) and included respiratory-related (occurring immediately following an apnea, hypopnea, or snore), technician-induced, and spontaneous arousals.

### *Behavior Apparatus and Procedures*

The Sleep Behavior Questionnaire (SBQ) provided support for the PSG results and was used to initially define conditions as control or snoring on the basis of parental reports. A proxy for IQ was obtained using the Peabody Picture Vocabulary Test–III (PPVT–III) and used as a screener. Participants' socioeconomic status and hand preferences (using the Edinburgh Handedness Inventory, Oldfield, 1971) were previously obtained and are not reported here. The Childhood Symptom Inventory–4 (CSI–4) screened for extreme behavioral disorders and characterized the neurocognitive status of children with and without SDB. The NEPSY is a standardized neuropsychological assess-

ment that has been used previously in studies of children with SDB. It was used as a measure of observable neurocognitive function. The Conners Parent Rating Scales–Revised (CPRS–R:L) is a widely used tool that is sensitive to the behavioral changes associated with SDB and habitual snoring. It was used as a measure of parental perception of neurobehavioral functioning and was the criterion predicted by the ERP variables in the multiple regression analysis described below.

### *Sleep Behavior Questionnaire*

Parents of all Sleep Medicine Center patients completed a detailed questionnaire about sleep problems, habits, and practices. Questions pertained to bedtime routines (e.g., “Will your child fall asleep alone in bed?” “In order to fall asleep does your child need a special toy or object?”), as well as sleep disturbances such as daytime sleepiness, restless sleep, nightmares, sleepwalking, nocturnal enuresis, sleep apnea, snoring, and nocturnal awakenings. The majority of questions were dichotomous with “yes” or “no” answers; however, some items were open-ended or multiple-choice responses (e.g., “What do you think that prevents your child from falling asleep?”). In previous studies from the Sleep Medicine Center, the SBQ had a high sensitivity and specificity for snoring and non-snoring children in this age range (Montgomery-Downs, O’Brien, Holbrook, & Gozal, 2004). For this study, the SBQ was used as a measure of parental report of sleep parameters, as the basis for group assignment as snoring or non-snoring, and to obtain background information such as demographics and history of sleep issues for each participant.

### *Childhood Symptom Inventory 4 (Gadow & Sprafkin, 2002)*

The CSI–4 Parent Checklist was used to screen for behavioral, emotional, and cognitive disorders based on parent-reported symptoms outlined in the *Diagnostic and Statistical Manual of Mental Disorders* (4th ed. [DSM–IV]; American Psychiatric Association, 2000). The tool was developed for use as a diagnostic and screening measure in clinical psychiatric and special education populations. Use of the CSI–4 to identify participants who should be excluded due to psychiatric or behavioral disorders was justified for this study. The questionnaire listed a variety of symptoms and the parent indicated whether the particular symptom “never,” “sometimes,” “often,” or “very often” characterized the child. The parent completed the CSI–4 at the initial overnight sleep study. If a child’s score indicated that he or she was at risk, a clinical psychologist evaluated the child. The internal consistency of the CSI–4 was high in general, with Cronbach’s alpha values ranging from .94 to .74 for the Parent Checklist categories. Test–retest reliability correlations for the symptom count and symptom severity scores were high (e.g., the Pearson Correlation for the AD/HD Hyperactive-Impulsive Type Symptom Severity Score between the initial session and the retest was 0.84) and all were significant at an alpha level of 0.0001 with a testing interval of 4 weeks. In this study, however, the CSI–4 was used only once (see aforementioned timeline) as a screening measure. The CSI–4 was used for children 5 to 8 years of age. Four-year-olds were given the version of the symptom inventory for younger children (Early Childhood Inventory 4, ECI–4). The diagnosis of a behavioral disorder was not determined from the CSI–4. Rather, children identified as “at-risk” due to a high symptom severity score on the CSI–4 (indicating a high occurrence of problem behaviors as identified by the *DSM–IV*) were referred to a clinical psychologist for a formal interview and assessment. The child was then excluded from the study if a clinical diagnosis was received.



*Peabody Picture Vocabulary Test—III (Dunn & Dunn, 1997)*

The PPVT—III was used as an IQ-screening tool and as a measure for characterizing the sample. The PPVT—III is the most widely used measure of receptive language abilities. It was co-normed with the Expressive Vocabulary Test thereby allowing for direct comparison of expressive and receptive language for individuals aged 2.5 to 90+ years. The assessment was individually administered and untimed, with children selecting the one picture of four choices that best corresponded to the word the tester read. The test has high reliability and internal consistency in young children and adults with reliability coefficients for the raw and standard scores ranging from 0.88 to 0.99 for all age groups (testing interval was one month) and Cronbach's alpha ranging from 0.92 to 0.98. For the purposes of this study, the PPVT—III was administered once. The assessment has high construct validity, with correlations between the PPVT—III standard scores and the Wechsler Intelligence Scale for Children—Third Edition (WISC—III) ranging from 0.82 to 0.92 for the norming sample. Vocabulary measures obtained with tests such as the PPVT—III are used as a proxy measure of overall intelligence and are related to language-based outcomes, such as those utilized in the present protocol. The test was administered before PSG and required approximately 15 to 20 min for children between 4 and 8 years of age.

*NEPSY (Korkman, Kemp, & Kirk, 1998)*

The NEPSY is a battery with five subtests used to provide an assessment of neuropsychological functioning in children from 3 to 12 years of age. These were measures of basic and complex cognitive capacities that affected learning and school performance. There are five domains: Attention/Executive Function, Language, Sensorimotor, Visuospatial Processing, and Memory and Learning. The NEPSY was administered twice as part of the larger overall study. Initial test scores for the Visual Attention subtest assessed whether ERP differences in response to an attention task relate to differences in observable behavior during a standardized attention test. During this subtest, children are shown an example of a picture ("bunny," "cat," or "faces") and asked to mark all identical figures in an array in under 3 min. Each child is given 2 arrays. The NEPSY has high test/retest reliability (e.g., an adjusted  $r$  squared of 0.9 for the Memory and Learning component) across sex and ages (3–12 years) even with a testing interval as short as 2 to 10 weeks. The internal consistency of the NEPSY is also high with  $r$  values ranging from 0.71 to 0.91 for all domains across the age range included in this study (4 to 8 years). The administration time for this test was approximately 6 min.

*Conners Parent Rating Scales-Revised: Long (Conners, 2000)*

The CPRS—R:L was originally published in 1970 by C. Keith Conners, PhD. The revised version was released in 1997. It was developed using a normative sample of over 8,000 children in the United States and Canada. It uses parental ratings to assess ADHD and problem behaviors in children aged 3 to 17 years. The CPRS—R:L has been widely used for research and clinical purposes, including screening, diagnosis, treatment planning, and outcome assessment. The questionnaire contains 80 Likert items assessing 7 factors: oppositional, cognitive problems/inattention, hyperactivity, anxious-shy, perfectionism, social problems, and psychosomatic. It provides 14 subscale scores: oppositional, cognitive problems/inattention, hyperactivity, anxious-shy, per-

fectionism, social problems, psychosomatic, ADHD Index, CGI: restless-impulsive, GCI: emotional lability, CGI: total, DSM-IV: inattentive, DSM-IV: hyperactive-impulsive, DSM-IV: total. The CPRS-R: L has been extensively used in other studies of pediatric SDB (O'Brien, Mervis, Holbrook, Bruner, Klaus, Rutherford, 2004b; Wei et al., 2007).

## ANALYSES

### Analysis of Electrophysiological Data

#### *Initial ERP Data Treatment*

Initial ERP data treatment or “preprocessing” was conducted using commercially available NetStation © software, version 4.1.2 (Electrical Geodesics, Inc., Eugene, OR). Single trial ERPs were screened following standard procedures for eye artifacts. Electrodes at standard electro-oculogram (EOG) positions measured eye movements. Trials with eye channel differences in excess of 70  $\mu\text{V}$  or more than 5% bad channels (defined as detecting voltage shifts in excess of 150  $\mu\text{V}$  within and across trials) were rejected. Channels characterized by consistent high voltages were replaced using a spherical interpolation algorithm. Following artifact screening, the EEGs were baseline corrected, using the 100 msec prestimulus average as a measure of baseline. ERPs were adjusted using a 30 Hz digital low pass filter and an average reference. Trials were then averaged for each subject separately (Andreassi, 2000; Regan, 1989).

#### *Principal Components Analysis (PCA)—Repeated Measures Analysis of Variance (RMA)*

Analysis of the preprocessed ERP data was conducted using the commercially available software package Statistical Software for the Social Sciences (SPSS) ©, version 11 for Mac OSX (SPSS, Inc., Chicago, IL). Investigators used spatial principal components analysis to identify variation in the distribution of activity across the scalp and determine regions of the topographic distribution that were significantly correlated. These covarying clusters of channels were then averaged to form the seven “virtual electrodes” outlined in Figure 3 in the Results section of this article. This reduced the data to meaningful measures and provided “virtual electrodes” that were subjected to temporal PCA with Varimax rotation to identify distinct regions of temporal variability in the waveform. The PCA-Varimax procedure was blind to individual experimental conditions and generated the same solution regardless of the order in which data were entered. PCA has less variance misallocation than traditional amplitude-latency measurements—or “peak-picking”—if sufficient power exists (Beauducel & Debener, 2003).

After PCA was used to identify the major ERP components, subsequent factor weights served as the dependent variables in a Repeated Measures Analysis of Variance (RMA) that tested the null hypothesis of  $\mu_{\text{SNORE}} = \mu_{\text{CONTROL}}$  for the electrophysiological task. The mixed-factorial RMA utilized the design, Group (2)  $\times$  Stimulus (2)  $\times$  Electrode Region (7), and identified significant interactions by group in the factor waveform data. Post-hoc corrected *t*-tests including Tukey's Honestly Significant Difference Test were performed when the omnibus RMA indicated significant interactions between group and within-subject variables such as stimulus (frequent or target) and electrode site. ERP variables that were significantly different between snoring children

and non-snoring controls were entered into a stepwise multiple regression to determine which ERP elements significantly predicted scores on the CPRS-R:L.

### *Source Localization*

Scalp potential source estimates were modeled using the GeoSource © imaging software (Electrical Geodesics, Inc., Eugene, OR). Investigators used a minimum-norm least squares solution (MNLS) and a finite difference model with the LAURA (local autoregressive average) constraint (Grave de Peralta Menendez et al., 2004) to derive source locations based on the Montreal Neurological Institute probabilistic magnetic resonance image (MRI). Settings included weighting per location, truncated singular value decomposition (TSVD) regularization at  $10^{-3}$ , and the radius of influence at 12.2 mm (Frishkoff, Perfetti, & Westbury, 2009). To compute estimates of the anatomical source activity, investigators focused on time points of maximal amplitude peaks from components of the temporal PCA. The source estimates contributing to the peaks for factors exhibiting significant differences by group are illustrated in the Results section.

### *Analysis of PSG Data*

Overnight sleep studies were performed on all participants to rule out sleep disorders and—for those individuals in the control group—confirm the absence of snoring. Investigators tested the null hypothesis that  $\mu_{\text{SNORE}} = \mu_{\text{CONTROL}}$  for sleep variables such as percent total sleep time (TST) in each sleep stage; number of respiratory, spontaneous, and total arousals; apnea/hypopnea index (AHI); and oxygen saturation using multiple un-paired Independent Samples *t*-tests.

### *Analysis of Behavioral Data*

Independent Samples *t*-tests were used to examine standardized scores from the screening instruments and demographic parameters to ensure there were no differences between groups in potential confounding variables such as body mass index (BMI) or estimated intelligence quotient (using the standardized score from the PPVT-III as a proxy). Investigators tested the null hypothesis that  $\mu_{\text{SNORE}} = \mu_{\text{CONTROL}}$  using standardized scores for the various domains of the CPRS-R:L and the Visual Attention subtest of the NEPSY. These served as the dependent variables for Independent Samples *t*-tests. Analyses were conducted to investigate significant differences between groups (Snoring versus Control). Variables on which the two groups were found to significantly differ were related to ERP data using stepwise multiple regression to determine if ERP differences could predict measures of behavior. An alpha level of 0.05 was used to determine significance for all statistical results including post-hoc and planned tests. Greenhouse-Geisser corrections were applied to all tests.

## RESULTS

### *ERP PCA and RMA Results*

Spatial PCA identified seven factors that accounted for 77.618% of the total variance in topographic distribution of activation. The resulting electrode groupings are illustrated in Figure 2.




FIGURE 2 Illustration of electrode groupings obtained from spatial principal components analysis.


Using those electrode groupings, the temporal PCA was performed, producing six temporal factors that accounted for 87.713% of the total variance in the model. The RMA was then performed using the factor weights from the temporal PCA as the dependent variable. Factors 1 and 5 were the only temporal factors to exhibit significant main effects of group or interactions by group.

The group\*stimulus interaction was significant for Factor 1 ( $F = 5.194, p = .028$ ) that peaked at 844 msec post-stimulus onset. Post hoc analyses indicated that there was no significant difference in the brain responses of snorers to target versus frequent stimuli. However, significant differences were noted between responses to stimuli in controls ( $t = -2.101, p = .043$ ). Additionally, there was a significant difference in how snoring versus control children processed the frequent tone at approximately 844 msec ( $t = -2.101, p = .042$ ).


There was a significant group\*electrode region interaction for Factor 5 ( $F = 4.915, p = .002$ ). Factor 5 had a peak at approximately 124 msec post-stimulus onset. Post hoc analyzes identified significant effects at two electrode regions. Control children exhibited increased ERP amplitude over electrode region 4, a centrally located region ( $t = 3.792, p = .001$ ). Snoring children exhibited increased ERP amplitude over electrode region 6, a right temporal region ( $t = -3.482, p = .001$ ). There were no significant differences between groups in the number of correct responses during the oddball task or in the reaction time for correct responses.

### Source Localization Results

Source localization solutions derived from topographical electrical activity and averaged by group are shown in Figures 3 and 4. In these topographic maps, one can see the statistically significantly increased activation at right temporal leads in snoring children compared to the more central activation recorded in non-snoring control children. Source localization was performed in GeoSource using this averaged brainwave data and produced the solutions illustrated in Figure 3. For snoring children, the estimated source generator in response to target stimuli was in the temporal lobe (specifically Brodmann area 22 in the superior temporal gyrus). The estimated source generator in response to the frequent stimuli was in the temporal lobe (Brodmann area 38 in the superior tem-



**FIGURE 3** Source localization by group at 124 msec (peak latency of temporal Factor 5). The sagittal view is on the left, coronal is in the center, and axial is on the right. For snoring children, the estimated source generators in response to target and frequent stimuli were in the temporal lobes. For control children, the estimated source generator in response to the target stimuli was located within the precuneus of the parietal lobe whereas the location shifted to the superior temporal gyrus of the temporal lobe in response to the frequent stimuli.



**FIGURE 4** Source localization by group at 844 msec (peak latency of temporal Factor 1). The sagittal view is on the left, coronal is in the center, and axial is on the right. In snorers, the estimated source generator in response to target and frequent stimuli was located in the frontal lobe. In controls, the estimated electrical source originated in the frontal lobe for target stimuli and in the amygdala for frequent stimuli.



poral gyrus). For control children, the estimated source generator in response to the target stimuli was in the precuneus of the parietal lobe (Brodmann area 31). The estimated source generator in response to the frequent stimuli for control children was located within the temporal lobe (Brodmann area 38 in the superior temporal gyrus).

Figure 4 illustrates the average activity for both groups at 844 msec post-stimulus onset, which is the latency of the peak of maximal activation for Factor 1. The significant difference in activation patterns for frequent versus target stimuli can be seen for children in the control group in the source localization. For snoring children, the estimated source generator in response to target stimuli was in the frontal lobe (specifically Brodmann area 11 in the orbital gyrus). The estimated source generator in response to the frequent stimuli was in the frontal lobe (Brodmann area 10 in the medial frontal gyrus). For control children, the estimated source generator in response to the target stimuli was in the frontal lobe (Brodmann area 10 in the medial frontal gyrus). The estimated source generator in response to the frequent stimuli for control children was in the limbic lobe (estimated in the uncus of the amygdala).

PSG Results

Group means for relevant parameters from the overnight polysomnographic studies are shown in Table 2. There were no significant differences by group in any clinically relevant markers for SDB, including apnea/hypopnea index or oxygen saturation nadir. There were, however, group differences in the percentage of total sleep time spent in sleep stages 2 and 4. Occasionally snoring children spent less time in stage 2 ( $t = -3.489, p = .001$ ) and more time in stage 4 ( $t = 2.040, p = .048$ ) than their non-snoring counterparts.

Behavioral and Descriptive Results

Descriptive and demographic information for the two groups is shown in Table 1. There were no significant differences between groups for any variable, including BMI. Mean standardized scores on the PPVT-III and scaled scores for the NEPSY Visual Attention subtest are also shown in

TABLE 2  
Polysomnographic Variables for Rarely Snoring and Non-Snoring Children

Variable	Snoring Group	Control Group
Apnea-Hypopnea Index	0.823 ± 0.866	0.850 ± 0.444
Apnea Index	0.627 ± 0.560	0.741 ± 0.386
Lowest Oxygen Saturation	93.619 ± 2.617	93.364 ± 2.682
% Stage 1	5.241 ± 2.725	4.923 ± 2.418
% Stage 2	45.427 ± 6.912*	52.086 ± 5.690*
% Stage 3	6.446 ± 2.823	5.686 ± 2.582
% Stage 4	22.064 ± 5.489**	18.686 ± 5.494**
% REM	19.841 ± 4.720	17.827 ± 4.178
Total Sleep Time	490.386 ± 39.490	480.318 ± 55.962
Arousal Index	8.88 ± 4.268	10.85 ± 5.415

Both groups have  $N = 22$ . Values given in means ± SD.  
\* $p < .05$ . \*\* $p < .01$ .

Table 1. There were no significant differences between groups on the PPVT–III, which was used as a proxy for IQ, or on the NEPSY Visual Attention scaled scores, which was used as a standardized assessment for attention. Rarely snoring children scored significantly higher than controls on 4 subscales of the CPRS–R:L: the Conners Global Index for restlessness and impulsivity ( $t = 2.314$ ,  $p = .028$ ); the cognitive problems/inattention subscale ( $t = 2.292$ ,  $p = .028$ ); the *DSM-IV*-based subscale for the ADHD inattentive subtype ( $t = 2.647$ ,  $p = .013$ ); and *DSM-IV*-based total index for the ADHD combined subtype ( $t = 2.402$ ,  $p = .023$ ).

The stepwise multiple regression analyses indicated that two ERP variables from Factor 1 significantly predicted scores on the CPRS–R:L. The factor weights corresponding to activation over electrode region 4 (including parietal electrode sites) significantly predicted scores on the cognitive problems/inattention subscale ( $F = 4.655$ ,  $p = .037$ ); the anxious/shy subscale ( $F = 4.586$ ,  $p = .038$ , adjusted R squared = 0.077); and the ADHD index ( $F = 4.098$ ,  $p = .049$ , adjusted R squared = 0.067). The factor weights corresponding to activation over electrode region 3 (including central and parietal electrode sites) significantly predicted CPRS–R:L scores on the perfectionism subscale ( $F = 5.131$ ,  $p = .029$ , adjusted R squared = 0.088).

## DISCUSSION

### ERP Results

Current ERP experiments show the presence of altered neural processing in children who snore occasionally and who otherwise have normal sleep studies. Indeed, the oddball task was more cognitively demanding for the rarely snoring children, who also exhibited more delayed and effortful processing. In contrast, control children processed the tones more quickly, engaging higher brain regions such as the precuneus at an earlier latency than snorers. At 124 msec post-stimulus, control children exhibited greater activation in attention-related central-parietal brain regions whereas snorers exhibited greater activation in the right temporal region, a region involved in pure tone discrimination. By 124 msec post stimulus, controls had already graduated to higher cognitive processes, while rare snorers were still engaged in basic sensory processing. Snoring children exhibited higher amplitude frontal activation than controls at 844 msec, indicating that once frontal processing occurred in rarely snoring children it required more effort than in control children. At this latency, children in the control group had less frontal activation and were showing activation in memory consolidation areas such as the amygdala—activation the rarely snoring children lacked.

Of note, no significant differences occurred between rarely snoring and non-snoring children on behaviorally observable phenomena such as the number of correct responses or the average reaction time for correct responses during the oddball task. It is our belief that rarely snoring children may be able to perform as well as the non-snoring children by engaging more cognitive resources (Molfese, Molfese, Beswick, Jacobi-Vessels, & Molfese, 2008). Our findings are consistent with previous reports in adults, suggesting that the volitional attention resources allocated to the processing of novel stimuli are sufficient in subjects with OSA, but more effortful and delayed relative to controls (Gosselin et al., 2006a; Gosselin et al., 2006b), most likely as the result of a decrease in overall cognitive resources available for the task at hand. Furthermore, the increased susceptibility of the prefrontal cortex and hippocampus to the effects of sleep fragmenta-

tion and hypoxia (D. Gozal, Daniel, & Dohanich, 2001; E. Gozal, 2001) associated with SDB could require reliance on suboptimal generators such as the cingulate, auditory, or parietal cortices (Gosselin et al., 2006b).

## Discussion of PSG Results

Analysis of PSG variables confirmed the initial hypothesis that rarely snoring children do not meet diagnostic criteria for clinically relevant SDB. The rarely snoring children in this study did not exhibit increased apnea or hypopnea frequencies than the controls. Notwithstanding, rarely snoring children experienced more behavioral problems than controls according to parental reports. This suggests that snoring—however infrequent—is linked to increased morbidity, and that current analytical methods used to score overnight sleep studies are not sufficiently sensitive to assess the overall impact of snoring on neuropsychological processes of children.

There was a somewhat unexpected finding upon analysis of PSG data. Rarely snoring children exhibited a decreased percentage of total sleep time in stage 2 of non-rapid eye movement (NREM) sleep and an increased percentage of total sleep time in stage 4 of NREM sleep. We are unclear as to the significance of such findings.

## Discussion of Behavioral Results

The hypothesis that snoring children would score significantly higher on subscales of the CPRS–R:L, thereby indicating increased behavioral problems, was also confirmed. Rarely snoring children scored higher on CPRS–R:L subscales for restlessness, cognitive problems/inattention, ADHD inattentive subtype, and ADHD combined subtype, similar to previous reports (Wei et al., 2007). Approximately one half of the snoring children scored one standard deviation or more above average on one or more subscales. In fact, most who scored above one standard deviation on one subscale scored high on several subscales.

Stepwise multiple regression findings supported the hypothesis that ERPs provide predictive markers for behavioral differences among the two study groups. For Factor 1, activity at electrode region 4 (including central and parietal electrode sites) elicited by target tones from the oddball task significantly predicted scores on the cognitive problems/inattention, anxious/shy, and ADHD indices of the CPRS–R:L. The parietal lobe contributes to nonverbal working memory (Wager & Smith, 2003), which could account for the accurate prediction of scores on the cognitive problems/inattention, perfectionism, and ADHD subscales. Activation at prefrontal-central electrode sites could explain the relationship between this electrode region and the anxious/shy subscale, which could incorporate internalizing or inhibiting behaviors. It is interesting to note that the group that showed no activation of the amygdala on source localization exhibited higher scores on the anxious/shy subscale. However, this difference was not statistically significant. The finding that two ERP variables can predict scores on subscales of the CPRS–R:L is particularly important given the general lack of bio-physiological assessments in psychological diagnostics, especially considering the great difficulty in assessing complex behaviors. The ability for a physiological assessment to account for nearly 10% of the variance of a behavioral assessment is not trivial. Such results underscore the argument that ERPs hold significant promise as quick, safe, and relatively inexpensive diagnostic tools.

### Study Limitations and Strengths

Reliance on parental report could have introduced bias. However, parental report of snoring is one of the most robust markers of neurocognitive impairment in studies of pediatric SDB (Beebe, 2006), suggesting the existence of a spectrum of snoring severity being associated with parallel changes in behavioral patterns. The lack of longitudinal follow-up to investigate long-term effects of non-habitual snoring in children is clearly a limitation of the present study and will require additional studies in the future. We did not incorporate analyses of genetic or proteomic markers such as Apolipoprotein E alleles or C reactive protein, which are altered in children with SDB, particularly in those who are most adversely affected by the underlying SDB (D. Gozal, Capdevila, et al., 2007; D. Gozal, Crabtree, Sans Capdevila, Witcher, & Kheirandish-Gozal, et al., 2007; Tauman, Ivanenko, O'Brien, & Gozal, 2004).

The strengths of this study are its prediction and explanation of real-life, behaviorally observed phenomena—such as the increased behavioral and attention problems previously reported in children with snoring and/or SDB—by objectively measured electrophysiological markers (Brietzke & Gallagher, 2006; O'Brien, Mervis, Holbrook, Bruner, Klaus, Rutherford, et al., 2004a; O'Brien, Mervis, Holbrook, Bruner, Smith, McNally, et al., 2004). Furthermore, there are many applications for this non-invasive and relatively inexpensive technique, which is well matched to the current need for new diagnostic and evaluative procedures in pediatric sleep medicine and research. Future studies are needed to link these results to brain morphology and more complex—and perhaps reciprocal—interactions between the brain, other systems such as the cardiovascular, metabolic, respiratory, and immune systems, and the complex neuropsychological process and behaviors that develop in childhood.

### CONCLUSIONS

Children who snore are not likely to “grow out of it” over the course of their development (Lofstrand-Tidestrom & Hultcrantz, 2007). In their 2008 review, Giordani and Chervin argue that SDB severity is not directly related to the degree of neurocognitive and behavioral impairment when assessed by polysomnography. For example, children with milder forms of SDB could have more behavior problems than children with overt OSA (Giordani & Chervin, 2008). The authors concluded that new measures should be developed and closely examined to further elucidate the relationship(s) between SDB and neurobehavioral morbidity. The present investigation represents one attempt in this direction.

Over 250,000 children are surgically treated for obstructive sleep apnea (the most severe form of SDB) each year in the United States alone (Brietzke & Gallagher, 2006). Evaluation and treatment of pediatric SDB is a major public health concern. Primary snoring, once considered important only insofar as it was a hallmark of more severe disease, is now being investigated as a milder form of SDB that may require treatment. Standard polysomnographic measures should no longer be considered as accurate predictors of neuropsychological morbidity in children who snore (Giordani & Chervin, 2008). Indeed, none of the children in this study met polysomnographic intervention criteria, and yet, substantial behavioral and ERP differences emerged when compared to control children. Rarely snoring children exhibited altered neural functioning, including delayed and more effortful processing, and a reliance on suboptimal cognitive generators. Thus, chil-

dren with a history of snoring of any frequency or severity should be excluded from any normative control group. More research is clearly needed to determine mechanisms underlying increased neurobehavioral susceptibility to the presence of snoring (Giordani & Chervin, 2008; D. Gozal, Capdevila, et al., 2007; Kuehni et al., 2008). Coupled with behavioral assessments, ERPs may prove useful in identifying which children would benefit from therapeutic interventions to reduce or eliminate snoring, and serve as useful tools to monitor treatment outcomes.

## REFERENCES

- Adams, A. S. M., Schluchter, M., Redline, S. (2001). Relation of measures of sleep-disordered breathing to neuropsychological functioning. *American Journal of Respiratory & Critical Care Medicine*, 163, 1626–1631.
- Alexopoulos, E. I., Kostadima, E., Pagonari, I., Zintzaras, E., Gourgoulialis, K., & Kaditis, A. G. (2006). Association between primary nocturnal enuresis and habitual snoring in children. *Urology*, 68(2), 406–409.
- Ali, N., Pitson, D., & Stradling, J. (1993). Snoring, sleep disturbance and behaviour in 4–5 year olds. *Archives of Disease in Childhood*, 68, 360–366.
- Ali, N. P., Pitson, D., & Stradling, J. R. (1996). Sleep disordered breathing: Effects of adenotonsillectomy on behaviour and psychological functioning. *European Journal of Pediatrics*, 155, 56–62.
- American Academy of Pediatrics, S. o. P. P., Subcommittee on Obstructive Sleep Apnea Syndrome. (2002). Clinical practice guideline: Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics*, 109, 704–711.
- American Sleep Disorders Association Task Force. (1992). *EEG arousals: Scoring rules and examples: A preliminary report from the Sleep Disorders Atlas Task Force of the American Sleep Disorders Association* (No. 0161-8105 (Print)).
- Andreassi, J. (2000). *Psychophysiology: Human behavior & physiological response* (Fourth ed.). Mahwah, NJ: Lawrence Erlbaum Associates.
- Arman, A. R., Ersu, R., Save, D., Karadag, B., Karaman, G., Karabekiroglu, K., et al. (2005). Symptoms of inattention and hyperactivity in children with habitual snoring: Evidence from a community-based study in Istanbul. *Child Care Health Development*, 31(6), 707–717.
- Ball, J., Tiernan, M., Janusz, J., & Furr, A. (1997). Sleep patterns among children with attention-deficit hyperactivity disorder: A re-examination of parental perceptions. *Journal of Pediatric Psychology*, 22, 389–398.
- Beauducel, A., & Debener, S. (2003). Misallocation of variance in event-related potentials: Simulation studies on the effects of test power, topography, and baseline to peak versus principal components qualifications. *Journal of Neuroscience Methods*, 124, 103–112.
- Beebe, D. (2006). Neurobehavioral morbidity associated with disordered breathing during sleep in children: A comprehensive review. *Sleep*, 29(9), 1115–1134.
- Beebe, D., & Gozal, D. (2002). Obstructive sleep apnea and the prefrontal cortex: Towards a comprehensive model linking nocturnal upper airway dysfunction to daytime cognitive and behavioral deficits. *Journal of Sleep Research*, 11, 1–16.
- Blunden, S., Lushington, K., Lorenzen, B., Martin, J., & Kennedy, D. (2005). Neuropsychological and psychosocial function in children with a history of snoring or behavioral sleep problems. *Journal of Pediatrics*, 146(6), 780–786.
- Brietzke, S. E., & Gallagher, D. (2006). The effectiveness of tonsillectomy and adenoidectomy in the treatment of pediatric obstructive sleep apnea/hypopnea syndrome: A meta-analysis. *Otolaryngology—Head & Neck Surgery*, 134(6), 979–984.
- Chervin, R. D., Ruzicka, D. L., Archbold, K. H., & Dillon, J. E. (2005). Snoring predicts hyperactivity four years later. *Sleep*, 28(7), 885–890.
- Conners, C. K. (2000). *Conners' rating scales—revised technical manual*. North Tonawanda, NY: Multi-Health Systems, Inc.
- Corkum, P., Tannock, R., Moldofsky, H., Hogg-Johnson, S., & Humphries, T. (2001). Actigraphy and parental ratings of sleep in children with attention deficit/hyperactivity disorder (ADHD). *Sleep*, 24, 303–312.
- Dunn, L., & Dunn, L. (1997). *Examiner's manual for the PPVT—III Peabody Picture Vocabulary Test—Third Edition*. Circle Pines, MN: American Guidance Service.
- Emmerson, R., Dustman, R., Shearer, D., & Turner, C. (1990). P3 latency and symbol digit performance correlations in aging. *Experimental Aging Research*, 15(3–4), 151–159.

- Frishkoff, G. A., Perfetti, C. A., & Westbury, C. (2009). ERP measures of partial semantic knowledge: Left temporal indices of skill differences and lexical quality. *Biological Psychology*, 80(1), 130–147.
- Gadow, K., & Sprafkin, J. (2002). *Childhood symptom inventory 4 screening and norms manual*. Stony Brook, NY: Checkmate Plus, LTD.
- Giordani, B., & Chervin, R. (2008). Sleep-disordered breathing and neurobehavioral outcomes: In search of clear markers for children at risk. *Journal of the American Medical Association*, 299(17), 2078–2080.
- Gislason, T., & Benediktsdottir, B. (1995). Snoring, apneic episodes, and nocturnal hypoxemia among children 6 months to 6-years-old. *Chest*, 107, 963–966.
- Gosselin, N., Mathieu, A., Mazza, S., Decary, A., Malo, J., & Montplaisir, J. (2006a). Deficits in involuntary attention switching in obstructive sleep apnea syndrome. *Neuroscience Letters*, 408, 73–78.
- Gosselin, N., Mathieu, A., Mazza, S., Petit, D., Malo, J., & Montplaisir, J. (2006b). Attentional deficits in patients with obstructive sleep apnea syndrome: An event-related potential study. *Clinical Neurophysiology*, 117(10), 2228–2235.
- Gottlieb, D., Vezina, R., Chase, C., Lesko, S., Heeeren, T., Weese-Meyer, D., et al. (2003). Symptoms of sleep-disordered breathing in 5-year-old children are associated with sleepiness and behavior problems. *Pediatrics*, 112, 870–877.
- Gozal, D. (1998). Sleep-disordered breathing and school performance in children. *Pediatrics*, 102, 616–620.
- Gozal, D., & Pope, D. (2001). Snoring during early childhood and academic performance at ages 13–14 years. *Pediatrics*, 107, 1394–1399.
- Gozal, D., Capdevila, O., Kheirandish-Gozal, L., & Crabtree, V. M. (2007). Apoe epsilon 4 allele, cognitive dysfunction, and obstructive sleep apnea in children. *Neurology*, 69(3), 243–249.
- Gozal, D., Crabtree, V. M., Sans Capdevila, O., Witcher, L. A., & Kheirandish-Gozal, L. (2007). C-reactive protein, obstructive sleep apnea, and cognitive dysfunction in school-aged children. *American Journal of Respiratory & Critical Care Medicine*, 176(2), 188–193.
- Gozal, D., Daniel, J., & Dohanich, G. (2001). Behavioral and anatomical correlates of chronic episodic hypoxia during sleep in the rat. *The Journal of Neuroscience*, 21(7), 2442–2450.
- Gozal, D., Holbrook, C., Mehl, R., Nichols, K. L., Raffield, T. J., Burnside, M. M., et al. (2001). Correlation analysis between NEPSY battery scores and respiratory disturbance index in snoring 6-year old children: A preliminary report. *Sleep*, 24, 126.
- Gozal, E., Row, B., Schurr, A., & Gozal, D. (2001). Developmental differences in cortical and hippocampal vulnerability to intermittent hypoxia in the rat. *Neuroscience Letters*, 305, 197–201.
- Grave de Peralta Menendez, R., Murray, M. M., Michel, C. M., Martuzzi, R., & Gonzalez Andino, S. L. (2004). Electrical neuroimaging based on biophysical constraints. *NeuroImage*, 21(2), 527–539.
- Guilleminault, C., Eldridge, F. L., Simmons, F. B., & Dement, W. C. (1976). Sleep apnea in eight children. *Pediatrics*, 58(1), 23–30.
- Guilleminault, C., & Lee, J. H. (2004). Does benign “Primary snoring” Ever exist in children? [comment]. *Chest*, 126(5), 1396–1398.
- Guilleminault, C., Winkle, R., Korobkin, R., & Simmons, B. (1982). Children and nocturnal snoring: Evaluation of the effects of sleep related respiratory resistive load and daytime functioning. *European Journal of Pediatrics*, 139(3), 165–171.
- Halbower, A. C., & Mahone, E. M. (2006). Neuropsychological morbidity linked to childhood sleep-disordered breathing. *Sleep Medicine Reviews*, 10(2), 97–107.
- Hill, W. (1889). On some causes of backwardness and stupidity in children: And the relief of the symptoms in some instances by nasopharyngeal scarifications. *British Medical Journal*, II, 711–712.
- Kaplan, B., McNichol, J., Conte, R., & Moghadam, H. (1987). Sleep disturbance in preschool-aged and hyperactive and nonhyperactive children. *Pediatrics*, 80, 839–844.
- Karpinski, A., Scullin, M., & Montgomery-Downs, H. (2008). Risk for sleep disordered breathing and executive function in preschoolers. *Sleep Medicine*, 9, 418–424.
- Key, A. D., Dove, G. O., & Maguire, M. J. (2005). Linking brainwaves to the brain: An ERP primer. *Developmental Neuropsychology*, 27(2), 183–215.
- Korkman, M., Kemp, S., & Kirk, U. (1998). *Nepsy: A developmental neuropsychological assessment*. San Antonio, TX: The Psychological Corporation.
- Kuehni, C. E., Strippoli, M. P., Chauliac, E. S., & Silverman, M. (2008). Snoring in preschool children: Prevalence, severity, and risk factors. *European Respiratory Journal*, 31(2), 326–333.
- Leach, J., Olson, J., Hermann, J., & Manning, S. (1992). Polysomnographic and clinical findings in children with obstructive sleep apnea. *Archives of Otolaryngology — Head & Neck Surgery*, 118, 741–744.



- Liukkonen, K., Virkkula, P., Aronen, E. T., Kirjavainen, T., & Pitkaranta, A. (2008). All snoring is not adenoids in young children. *International Journal of Pediatric Otorhinolaryngology*, 72(6), 879–884.
- Lofstrand-Tidestrom, B., & Hultcrantz, E. (2007). The development of snoring and sleep related breathing distress from 4 to 6 years in a cohort of Swedish children. *International Journal of Pediatric Otorhinolaryngology*, 71(7), 1025–1033.
- Lopes, M. C., & Guilleminault, C. (2006). Chronic snoring and sleep in children: A demonstration of sleep disruption. *Pediatrics*, 118(3), e741–746.
- Lyytinen, H., Ahonen, T., Eklund, K., Guttorm, T. K., Laakso, M. L., Leinonen, S., et al. (2001). Developmental pathways of children with and without familial risk for dyslexia during the first years of life. *Developmental Neuropsychology*, 20(2), 535–554.
- Lyytinen, H., Blomberg, A. P., & Naatanen, R. (1992). Event-related potentials and autonomic responses to a change in unattended auditory stimuli. *Psychophysiology*, 29(5), 523–534.
- Lyytinen, H., & Naatanen, R. (1987). Autonomic and erp responses to deviant stimuli: Analysis of covariation. *Electroencephalography & Clinical Neurophysiology - Supplement*, 40, 108–117.
- Mitchell, R. B., & Kelly, J. (2005). Child behavior after adenotonsillectomy for obstructive sleep apnea syndrome. *Laryngoscope*, 115(11), 2051–2055.
- Molfese, D. L. (1989). The use of auditory evoked responses recorded from newborn infants to predict later language skills. *Birth Defects: Original Article Series*, 25(6), 47–62.
- Molfese, D. L. (1990). Auditory evoked responses recorded from 16-month-old human infants to words they did and did not know. *Brain & Language*, 38(3), 345–363.
- Molfese, D. L. (2000). Predicting dyslexia at 8 years of age using neonatal brain responses. *Brain & Language*, 72(3), 238–245.
- Molfese, D. L., & Molfese, V. J. (1985). Electrophysiological indices of auditory discrimination in newborn infants: The bases for predicting later language development. *Infant Behavior and Development*, 8, 197–211.
- Molfese, D. L., Key, A. F., Kelly, S., Cunningham, N., Terrell, S., Ferguson, M., et al. (2006). Below-average, average, and above-average readers engage different and similar brain regions while reading. *Journal of Learning Disabilities*, 39(4), 352–363.
- Molfese, D., Molfese, V., Beswick, J., Jacobi-Vessels, J., & Molfese, P. (2008). Dynamic links between emerging cognitive skills and brain processes. Special issue on neurobiological and experiential dimensions of dyslexia: Multiple perspectives. *Developmental Neuropsychology*, 33(6), 682–706.
- Montgomery-Downs, H. E., O'Brien, L., Gulliver, T., & Gozal, D. (2006). Polysomnographic characteristics in normal preschool and early school-aged children. *Pediatrics*, 117(3), 741–753.
- Montgomery-Downs, H. E., O'Brien, L., Holbrook, C. R., & Gozal, D. (2004). Snoring and sleep-disordered breathing in young children: Subjective and objective correlates. *Sleep*, 27(1), 87–94.
- O'Brien, L., Mervis, C., Holbrook, C., Bruner, J., Klaus, C., Rutherford, J., et al. (2004a). Neurobehavioral implications of habitual snoring in children. *Pediatrics*, 114(1), 44–49.
- O'Brien, L., Mervis, C., Holbrook, C., Bruner, J., Smith, N., McNally, N., et al. (2004). Neurobehavioral correlates of sleep-disordered breathing in children. *Journal of Sleep Research*, 13(2), 165–172.
- O'Brien, L., Mervis, C. B., Holbrook, C. R., Bruner, J. L., Klaus, C. J., Rutherford, J., et al. (2004b). Neurobehavioral implications of habitual snoring in children. *Pediatrics*, 114(1), 44–49.
- O'Brien, L., Tauman, R., & Gozal, D. (2004). Sleep pressure correlates of cognitive and behavioral morbidity in snoring children. *Sleep*, 27(2), 279–282.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh Inventory. *Neuropsychologia*, 9, 97–113.
- Owens, J., Maxim, R., Nobile, C., McGuinn, M., & Msall, M. (2000). Parental and self-report of sleep in children with attention deficit/hyperactivity disorder. *Archives of Pediatrics & Adolescent Medicine*, 154, 549–555.
- Polich, J., & Martin, S. (1992). P300, cognitive capability, and personality: A correlational study of university undergraduates. *Personality and Individual Differences*, 13, 533–543.
- Rechtschaffen, A., & Kales, A. (1968). *A manual of standardized terminology: Techniques and scoring system for sleep stages of human subjects*. Washington, DC: US Public Health Service.
- Regan, D. (1989). *Human brain electrophysiology: Evoked potentials and evoked magnetic fields on science and medicine*. New York: Elsevier Science Publishing, Inc.
- Rhodes, S., Shimoda, K., Wald, L., O'Neil, P., Oexmann, M., Collop, N., et al. (1995). Neurocognitive deficits in morbidly obese children with obstructive sleep apnea. *Journal of Pediatrics*, 127(741–744).

- Ring, A., Stein, D., Baiak, Y., Teicher, A., Hadjez, J., Elizur, A., et al. (1998). Sleep disturbances in children with attention-deficit/hyperactivity disorder: A comparative study with healthy siblings. *Journal of Learning Disabilities*, 31, 572–578.
- Rosen, C., Storfer-Isser, A., Taylor, H., Kirchner, H., Emancipator, J., & Redline, S. (2004). Increased behavioral morbidity in school-aged children with sleep-disordered breathing. *Pediatrics*, 114(6), 1640–1648.
- Singer, L. S. P. (1990). Complications of pediatric obstructive sleep apnea. *Otolaryngologic Clinics of North America*, 23, 665–676.
- Society, A. T. (1996). Standards and indications for cardiopulmonary sleep studies in children. *American Journal of Respiratory & Critical Care Medicine*, 153, 866–878.
- Stein, M. (1999). Unraveling sleep problems in treated and untreated children with adhd. *Journal of Child and Adolescent Psychopharmacology*, 9, 157–168.
- Sutton, S., Tuetting, P., Zubin, J., & John, E. R. (1965). Evoked potential correlates of stimulus uncertainty. *Science*, 150, 1187–1188.
- Tauman, R., Ivanenko, A., O'Brien, L., & Gozal, D. (2004). Plasma C-reactive protein in children with sleep-disordered breathing. *Pediatrics*, 113, 564.
- Urschitz, M., Guenther, A., Eitner, S., Urschitz-Duprat, P., Schlaud, M., Ipsiroglu, O., et al. (2004). Risk factors and natural history of habitual snoring. *Chest*, 126, 790–800.
- Urschitz, M. S., Guenther, A., Eggebrecht, E., Wolff, J., Urschitz-Duprat, P. M., Schlaud, M., et al. (2003). Snoring, intermittent hypoxia and academic performance in primary school children. *American Journal of Respiratory & Critical Care Medicine*, 168(4), 464–468.
- Wager, T., & Smith, E. (2003). Neuroimaging studies of working memory: A meta analysis. *Cognitive, Affective, & Behavioral Neuroscience*, 3(4), 255–274.
- Wei, J., Mayo, M., Smith, H., Reese, M., & Weatherly, R. (2007). Improved behavior and sleep after adenotonsillectomy in children with sleep-disordered breathing. *Archives of Otolaryngology and Head and Neck Surgery*, 133(10), 974–979.
- Weissbluth, M. D. A., Poncher, J., Reiff, J. (1983). Signs of airway obstruction during sleep and behavioral, developmental and academic problems. *Journal of Developmental & Behavioral Pediatrics*, 4, 119–121.